



## Chemical analysis of novel ginger and *Hibiscus sabdariffa* bio-preserved tomato powder

Olatayo A. Afolabi<sup>1</sup>, Omowumi T. Kayode<sup>1\*</sup>, Abolanle A. Kayode<sup>2</sup> and Gabriel O. Ajayi<sup>1</sup>

<sup>1</sup>Department of Biochemistry, College of Basic and Applied Sciences, Mountain Top University, Ogun State, Nigeria

<sup>2</sup>Department of Biochemistry, School of Basic Medical Sciences, Babcock University, Ilishan-Remo, Ogun State, Nigeria

\*Corresponding author: Kayode O.T., Ph.D., Department of Biochemistry, College of Basic and Applied Sciences, Mountain Top University, Ogun State, Nigeria.

**Submission Date:** July 8<sup>th</sup>, 2023; **Acceptance Date:** August 4<sup>th</sup>, 2023; **Publication Date:** August 11<sup>th</sup>, 2023

**Please cite this article as:** Afolabi O. A., Kayode O. T., Kayode A. A., Ajayi G. O. Chemical analysis of novel ginger and *Hibiscus sabdariffa* bio-preserved tomato powder. Bioactive Compounds in Health and Disease 2023; 6(8): 172-184. DOI: <https://www.doi.org/10.31989/bchd.v6i8.1129>

### ABSTRACT

**Background:** Tomatoes (*Solanum lycopersicum* L.) are widely consumed vegetables globally and are well known for their nutrients and economic value. This research aimed to create a tomato powder with improved nutraceutical benefits based on its ingredients.


**Methods:** Fresh tomatoes, ginger, and dried hibiscus calyx were obtained and dehydrated at 45 °C for 10 hours, 8 hours, and 1 hour and 30 minutes respectively. Proximate, elemental, phytochemical, High-performance Liquid Chromatography (HPLC), and Gas Chromatography-Mass Spectrometry (GCMS) analysis of the tomato powder mix were determined using standard methods.

**Results:** The proximate analysis revealed constituents of carbohydrates, fats, proteins, ash, fiber, and moisture as 65, 5, 10, 1, 7, and 12 % respectively. The GCMS also revealed 36 classes of compounds with different retention times. The most prominent peak is oleic acid with an RT value of 20.9. The fiber content of the tomato powder mix was higher than what is obtainable in fresh tomatoes. Bioactive compounds like stigmasterol (5.25%), oleic acids (49.11%), n-hexadecenoic acid (26.58%), campesterol (2.72%), and octadecanoic acid (6.8%) are notably present with lycopene (135.745 mg/100 grams).

**Conclusion:** The tomato powder mix is nutrient-packed and non-toxic. The bioactive components are known as

antioxidant, anti-inflammatory, cardiovascular health enhancers, etc showing the possible use of the powder as a functional food.

**Keywords:** Tomatoes powder, bioactive compounds, functional foods, dehydration



©FFC 2023. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0>)

## INTRODUCTION

Tomato (*Solanum lycopersicum* L.) is a widely grown and consumed vegetable globally due to its high economic viability, culinary flavors and texture, and most especially its health benefits. Modal described functional foods as any natural or refined food that has health benefits and disease prevention activities above their simple nutritional value [1]. Tomatoes have been classified as a functional food because of

their anti-inflammatory, anti-atherogenic, antioxidant, and anti-cancer properties. They also lower the risk of cardiovascular disease [2,4]. It is used for making various delicacies and sauces, soups, and salads; processed into ketchup, tomato pastes, tomato puree, powdered, flaked, whole peeled tomatoes, and canned, and diced tomatoes [5,6]. The climacteric nature of tomatoes accounts for the high post-harvest losses synonymous with its production

globally especially in developing countries, despite the annual increase in production. Dehydration is one of the oldest, most convenient, and widely used methods for preserving tomatoes. Numerous benefits of tomato powder include ease of drying, packaging, transportation, and storage. It is in high demand currently due to its use in making convenient and fast foods like pizzas both locally and internationally [7,8]. However, dehydration also comes with the challenge of reduction in the concentration of bioactive components, loss of color, and unwanted taste. Hence, there is a need to have a quality enhanced dehydrated tomato using other functional foods. Ginger (*Zingiber officinale* R) is one of the most popular nutritional condiments in the world with a distinctive pungent flavor loaded with bioactive compounds [9]. The rhizome of the plant is consumed and dated back in Chinese history for its therapeutic properties in preventing and managing arthritis, migraines, colds, nausea, and hypertension [10,12]. *Hibiscus sabdariffa* Linn. is a traditional plant used for medicine, herbal drinks, and delicacies due to its high phytochemical content. The leaves are eaten as vegetables, the calyces are used to make herbal drinks and medications, and the buds are used to make jam and preserves [13]. It has been shown to have antibacterial, antioxidant, anti-cholesterol, anti-diabetic, anti-hypertensive, anti-obesity, anti-anemia, nephron- and hepato-protective diuretic effects which are responsible for its use in the treatment of hypertension and hyperlipidemia. The bioactive molecules found in roselle are primarily responsible

for their positive effects on health [14,15].

The objective of this research was to produce a tomato powder made by dehydration with high bioactive components, good organoleptic taste, pleasant physical appearance and enhanced bioactive compounds without artificial additives to reduce wastage and increase food security.

## METHODS

**Powder Preparation:** Fresh red and ripened tomatoes weighing 800 grams, and yellowish–brown fresh ginger roots weighing 240 grams without any bruises were selected from the bulk buying from the Mile 12 market in Lagos, Nigeria. Dried reddish–black hibiscus calyx weighing 60 grams was also obtained from the same market.

The fresh fruits were sanitized in 1 % NaCl solution for 2 minutes and thoroughly rinsed with clean tap water and dried with a clean kitchen towel. The tomatoes were sliced to a very thin thickness using a sharp kitchen knife. The slices were vacuum dried at 45 °C for 10 hours, milled in a high-speed kitchen blender, and sieved through a 500 micrometer (µm) aperture size sieve for even powder, packed in air-tight ziplock pouches and stored under normal conditions. The cleaned ginger roots were peeled and cut into thin slices using a sharp kitchen knife. The slices were vacuum dried at 45 °C for 8 hours, milled in a high-speed kitchen blender, and sieved through a 500 micrometer (µm) aperture size sieve for even powder, packed in airtight Ziplock pouches and stored under normal conditions. The dried hibiscus calyx was hand-picked to get a

homogenous sample. It was rinsed briefly with clean water to remove all forms of dust, drained, and dehydrated at 40 °C for 1 and half hours (until no further difference in weight was observed), milled in a high-speed kitchen blender, and sieved through a 500 micrometer ( $\mu\text{m}$ ) aperture size sieve for even powder, packed in airtight Ziplock pouches and stored under normal conditions. The tomato powder mix with the best output (tomatoes, ginger, and hibiscus in the ratio of 5:2:1) was used for further analysis.

**Proximate analysis:** These analyses were carried out according to the method described by Kayode *et al* [16]. The moisture level was determined at 100 °C for 2 hours in an oven (no further weight loss was observed). The ash was determined after 2 hours at 600 °C in the furnace. The crude fiber value was determined according to Weende's method. The crude protein was calculated from the total nitrogen value obtained by multiplying it with a 6.25 factor using the Kjeldahl method. The crude fat was extracted by the Soxhlet method at 40-60 °C for 4 hours and the carbohydrate was determined by the differential method using mathematical calculation as nitrogen-free extract (NFE) which represents the soluble carbohydrate and other digestible and easily utilizable non-nitrogenous substances.  $\% \text{NFE} = 100 - (\% \text{moisture} + \% \text{crude fiber} + \% \text{crude protein} + \% \text{crude fat} + \% \text{Ash})$  [17].

**Mineral elements determination:** This was carried out after the digestion, ashing, and filtration of the

powder mix in 1 N Nitric acid  $\text{HNO}_3$ , 600 °C, and 1 N HCL respectively. Presence and concentration of Fe, Al, Mn, Pb, Ni, Cu, Zn, Se, Co, Cd, Ba, Cr, Va, As, and Hg. Pb, Cd, and Ni were determined by anhydride generation- HGA graphite furnace using high purity argon while other measurements were carried out in an air/acetylene flame Atomic-Absorption (AAS) (Perkin Elmer AAnalyst 700 model AAS) with a deuterium background corrector [16,18].

**Phytochemical analysis:** Qualitative analysis of Alkaloids, anthocyanin, flavonoids, phenols, glycosides, terpenoids, steroids, and phlobatannins was carried out following Banu and Cathrine [19,21]. One mL of ethanol and 1 mL of 10 % ferric chloride were combined with about 0.1 g of the extract. The presence of flavonoids was indicated by a brown solution with a dirty green precipitate [20].

The ferric chloride test was also used to confirm the presence of tannin and phenol compounds in the tomato mix sample. Five milliliters (5 mL) of distilled water were added to 50 mg of the powder mix. Five percent of neutral ferric chloride was added to the sample solution. The presence of tannin and phenol was confirmed by the green color [19].

Borntrager's test was used to determine the presence of glycosides and anthraquinones in the tomato powder mix. The powder mix was hydrolyzed (50 mg) with concentrated hydrochloric acid for 2 hours in a water bath. Two milliliters (2 mL) of the filtered hydroxylate were mixed with 3 mL of

chloroform and shaken. 10% ammonium solution was added after the chloroform layer was separated. The pink color indicated the presence of glycosides and anthraquinone in the upper aqueous layer [22].

Alkaloids was determined using Mayer's test. 5 mg of the powder mix was dissolved in 5 mL of distilled water. Two drops of Mayer's reagent were added to 3 mL of the sample through the sides of the test tube. White precipitate confirms the presence of alkaloids [22].

**Lycopene determination:** This was carried out using HPLC with slight modifications to Anguelova and Warthesen [23]; and Barba *et al* [24] methods. Tetrahydrofuran and methanol (1:1 v/v THF: MeOH) were used to extract the lycopene from the sample. The sample (0.2 mL) was reconstituted by adding 10 mL of distilled water, vortexed, and 40 mL of THF: MeOH mixture was added before centrifuging at 5000 rpm for 5 minutes and the supernatant was collected. This procedure was repeated several times until a colorless supernatant was obtained. Under nitrogen, the supernatant was evaporated to dryness and the residue redissolved in 4 mL hexane, filtered (0.45  $\mu$ m), and 100  $\mu$ l injected into the HPLC. Reverse phase HPLC was carried out on a C18 (201TP540) analytical column (5  $\mu$ m, 25 cm 4.6 mm; VYDAC, Hesperia, Calif., USA) with absorbance at 475 nm, flow rate of 1 mL/minute, and column temperature of 30°C.

Bioactive components of the powder mix were determined using a spectrophotometer GC-MS Agilent 7890A, GC/5975C MS (Agilent Technologies, Santa Clara,

USA according to the method described by Tang *et al* [25]. In a glass vial, 400 mg of the sample, 2 mL of methanol, and 20  $\mu$ l of internal standard ribitol were mixed and vortexed for 10 seconds, ultrasonicated for 30 minutes at 70 °C and centrifuged at 12,000 g for 10 minutes. One ml of chloroform was added to the supernatant with an equal volume of water and the mixture was centrifuged at 14000 g for 5 minutes and 400  $\mu$ l of the supernatant dried with a nitrogen-blowing device. This was followed by the addition of 80  $\mu$ l of methoxamine hydrochloride solution (15 g L<sup>-1</sup>, in pyridine) to the dried sample, vortexed for 30 s, and incubated at 37 °C for 90 minutes. Lastly, 80  $\mu$ l of Bis trimethylsilyl trifluoroacetamide (BSTFA) with 1 % trimethylchlorosilane (TMCS) was added to trimethylsilylation and heated for 1 hour at 70 °C. The mixture was then analysed using GC-MS.

**Statistical analysis:** Data is expressed in percentage from mean  $\pm$  SEM (standard error mean) in triplicates.

## RESULTS

The proximate analysis of the tomato powder mix showed a high carbohydrate content, good fiber, and reduced moisture content (Figure 1). The mineral analysis data (Table 1) showed low concentrations of heavy metals, Ag, Pb, Ni, Se, and As were all present below the limit of quantification (LOQ). Manganese, Fe, and Al are present in moderate concentration. The phytochemical screening confirmed the presence of alkaloids, tannins, steroids, flavonoids, glycosides, anthocyanins, anthraquinones, terpenes, and coumarins

Proximate analysis:




Figure 1: Proximate analysis of tomato powder mix

Table 1: Mineral elements determination

Minerals	Concentration (ppm)	LOD	LOQ	Correlation coefficient
Aluminium	2.0132	0.2532	0.7671	0.9991
Arsenic	<LOQ	0.2056	0.6230	0.9995
Cadmium	<LOQ	0.0357	0.1081	0.9996
Cobalt	<LOQ	0.3260	0.9878	0.9990
Chromium	<LOQ	0.1657	0.5022	0.9996
Copper	<LOQ	0.7679	0.5087	0.9996
Iron	1.1309	0.1451	0.4397	0.9997
Manganese	1.3566	0.2584	0.7830	0.9991
Nickel	<LOQ	0.2101	0.6366	0.9994
Lead	<LOQ	0.1969	0.5965	0.9994
Selenium	<LOQ	1.1739	3.5572	0.9825
Vanadium	<LOQ	0.1821	0.5520	0.9995
Zinc	<LOQ	0.1978	0.5994	0.9995
Silver	<LOQ	0.2621	0.7944	0.9991
Barium	<LOQ	0.2702	0.8188	0.9995

**Qualitative phytochemicals screening of tomato**

**powder Mix:** The quantity of lycopene in 100 g of the sample is shown in Table 2. The bioactive component data is presented in Table 3. Oleic acid has the highest

concentration, hexadecenoic acid, stigmasterol, octadecanoic acid, and campesterol are shown to be present and heptadecanoic acid is the least.

**Quantitative phytochemicals screening of tomato powder mix:****Table 2:** Lycopene quantity in tomato powder mix as identified by the HPLC.

Area	RETENTION TIME	LIBRARY ID	MOLECULAR WEIGHT	MOLECULAR FORMULA	Amount mg/100g
5.250	19.698	Lycopene	538.9	C <sub>40</sub> H <sub>56</sub>	135.745

**Bioactive components of tomato powder mix using GC-MS:****Table 3:** Chemical components in tomato powder mix as identified by GC-MS.

PEAKS	RT	LIBRARY ID	MOLE. WEIGHT	MOLE. FORMULA	AMOUNT mg/100g
1	3.39	Mesitylene	120.19	C <sub>9</sub> H <sub>12</sub>	0.17
2	3.55	2-Heptafluorobutyroxydedocane	382.36	C <sub>16</sub> H <sub>25</sub> F <sub>7</sub> O <sub>2</sub>	0.10
3	3.76	Benzene, 1-methyl-3-propyl	202.22	C <sub>11</sub> H <sub>13</sub> F <sub>3</sub>	0.10
4	4.44	Undecane	156.31	C <sub>11</sub> H <sub>24</sub>	0.23
5	4.47	1,3-cyclopentadiene, 1,2,3,4-tetramethyl-5-methylene	392.80	C <sub>16</sub> H <sub>32</sub> P <sub>2</sub> Pd <sup>-2</sup>	0.11
6	12.23	E-Dodec-2-en-1-yl propyl carbonate	270.41	C <sub>16</sub> H <sub>30</sub> O <sub>3</sub>	0.10
7	12.84	Tetradecanoic acid	228.37	C <sub>14</sub> H <sub>28</sub> O <sub>2</sub>	0.29
8	13.13	Octadecane	290.9	C <sub>18</sub> H <sub>37</sub> Cl	0.13
9	13.67	Hexahydropyridine 1-methyl-4-[4,5-dihydroxyphenyl]	207.27	C <sub>12</sub> H <sub>17</sub> NO <sub>2</sub>	0.16
10	13.86	Pentadecanoic acid	242.4	C <sub>15</sub> H <sub>30</sub> O <sub>2</sub>	0.37
11	14.69	Methyl hexadic-9-enoate	268.4	C <sub>17</sub> H <sub>32</sub> O <sub>2</sub>	0.44
12	26.58	Palmitic acid	256.42	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	26.58
13	15.61	9-cycloheptadecen-1-one	250.4	C <sub>17</sub> H <sub>30</sub> O	0.38
14	15.82	Margaric acid	270.5	C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	0.39
15	15.92	Ergost-4,7,22-trien-3, alpha-ol	396.6	C <sub>28</sub> H <sub>44</sub> O	0.17
16	16.01	2-hydroxychalcone	224.25	C <sub>15</sub> H <sub>12</sub> O <sub>2</sub>	0.14
17	16.07	Stigmasta-4, 22-diene	410.70	C <sub>29</sub> H <sub>46</sub> O	0.20
18	16.26	Ergosta-4,6,22-trien-3, beta-ol	396.6	C <sub>28</sub> H <sub>44</sub> O	0.39

PEAKS	RT	LIBRARY ID	MOLE. WEIGHT	MOLE. FORMULA	AMOUNT mg/100g
19	16.66	E-Oleic acid	49.11	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	49.11
20	16.79	Stearic acid	284.5	C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	6.80
21	17.45	4-methyl-1,4-heptadiene	110.2	C <sub>8</sub> H <sub>14</sub>	0.26
22	18.10	E-E-10,12-hexadecadien-1-ol acetate	280.4	C <sub>18</sub> H <sub>32</sub> O <sub>2</sub>	0.40
23	18.22	Z-Z-10,12-Hexadecadien-1-ol acetate	280.4	C <sub>18</sub> H <sub>32</sub> O <sub>2</sub>	0.38
24		2(1H)-Naphthalone,(octahydro-4a-methyl-7-(1-methylethyl),(4a alpha, 7 beta, 8a beta)	208.34	C <sub>14</sub> H <sub>24</sub> O	0.16
25	18.43	2(1H)-Naphthalone,(octahydro-4a-methyl-7-(1-methylethyl),(4a alpha, 7 beta, 8a beta)	208.34	C <sub>14</sub> H <sub>24</sub> O	0.80
26	18.66	Cyclodecanol	184.32	C <sub>12</sub> H <sub>24</sub> O	0.18
27	18.87	9,17-Octadecadienal	264.4	C <sub>18</sub> H <sub>32</sub> O	0.24
28	19.30	Naphthalene 1,2,3,4-tetrahydro-5-nitro	177.2	C <sub>10</sub> H <sub>11</sub> NO <sub>2</sub>	0.27
29	19.66	1,22-docosanediol	342.60	C <sub>22</sub> H <sub>46</sub> O <sub>2</sub>	0.35
30	19.96	Campesterol	400.7	C <sub>28</sub> H <sub>48</sub> O	2.72
31	10.25	Dihydrotachysterol	398.7	C <sub>28</sub> H <sub>46</sub> O	0.68
32	20.44	1-heptadecanamine	283.5	C <sub>19</sub> H <sub>41</sub> N	0.14
33	20.67	1H-Indene 2-butyl-5-hexyloctahydro	264.5	C <sub>19</sub> H <sub>36</sub>	0.35
34	20.93	Stigmasterol	412.7	C <sub>29</sub> H <sub>48</sub> O	5.25
35	21.20	Stigmasta-3,5-diene	396.7	C <sub>29</sub> H <sub>48</sub>	0.71
36	21.61	Cholesta-3,5-diene	368.6	C <sub>27</sub> H <sub>44</sub>	0.765

## DISCUSSION

**Proximate analysis:** Fresh tomatoes have a moisture content range of 89 to 93 %, which according to Aboagye Nuamah et al [26], is responsible for their susceptibility to microbial attack and results in their shorter shelf life. A longer shelf life than fresh tomatoes is indicated by the 12 % moisture in the powder mixture. The results from Obadina *et al.*, and others [27-29], gave higher moisture content although Degwale et al. [30] showed similar and lower results but at higher temperatures. Fibre is a component of food that

promotes weight loss, normalizes blood lipid levels, lowers the risk of cardiovascular diseases, promotes gastrointestinal health, and lowers the risk of certain diseases like certain cancers (colon cancer) and constipation [31,32] The fiber concentration of this study was higher than the values previously reported by Islay *et al.* [33], Bello and Amubieva [34], and Sanusi *et al.* [17] because ginger and roselle were added to the powder. Thus, it can be used as a functional food ingredient to improve overall health and possibly aid in weight loss. In finished product, acidity enhances flavor



and increases food safety by preventing the growth of microorganisms (reducing spoilage). The acidity is found to be within permissible limits [26,27].

**Minerals:** These are essential elements that cannot be synthesized by cells and are required in large or in minute quantities for cellular functions [35]. The mineral content determination of the powder showed that varied minerals quantities were present, and their levels were within approved daily nutritional requirements. A variety of trace minerals are involved in the regulation of protein synthesis and enzyme activities involved in cellular antioxidation and inflammation [36]. The composition of these minerals in the tomato-mix powder will improve metabolic function when ingested regularly, as well as improve the redox status of consumers which can help ameliorate myriads of metabolic diseases.

**Secondary metabolites:** These are known as bioactive compounds in foods and are either nutrients or non-nutrients that are found in small amounts in food and are associated with a healthy state beyond their basic nutritional values [37]. Lycopene, a crucial carotenoid in tomatoes, has been found to regulate anticancer, antioxidant, and anti-inflammatory effects associated with tomato consumption in studies comparing tomatoes and pure lycopene in reducing induced inflammation [38]. Lycopene acts as a chain-breaking antioxidant with a greater ability to scavenge singlet oxygen, superoxide hydroxyl, and peroxy than other antioxidants [39], eating tomatoes most likely decreases the risk of cardiovascular diseases and the atherogenic index [40,41]. The lycopene content (135.74 mg/100 g) in the tomato powder mix

revealed a higher level compared to those previously reported [42,43] This may be as a result of *Hibiscus sabdarifa* which is found to contain lycopene and some other carotenoids though in lower concentration compared to tomatoes [44]. The constituents of the tomato powder mix altered the bioactive compounds present when compared to pure tomato powder. According to Yusop et al [45,46], thirty-one compounds were detected using the methanol-chloroform extraction method compared to the 36 peaks detected in the tomato powder mix. The tomato powder mix showed a higher concentration of bioactive compounds like oleic acid and stigmasterol [47]. Oleic acid is a monounsaturated fatty acid which is also known as omega-9, it can reduce cholesterol and the risk of developing heart disease. The high oleic acid in this work suggests that the tomato powder mixture is very nutritive in addition to its anti-inflammatory, antioxidant, and anticancer properties [48-50]. According to a previous report, roselle has a high (28%) oleic acid [51] which may be responsible for the higher value in the powder mix. According to Bohm [41], oleic acid and lycopene have already been formulated into products for treating chronic inflammation, cancer, and cardiovascular diseases in America and Europe.

Stigmasterol is an unsaturated phytosterol that appears in plants as fats or oils and is present in vegetables, nuts, and seeds. It is well known to have anti-inflammatory, anti-cancer, anti-tumor, antioxidant, anti-osteoarthritic, anti-hypercholesterolemic, and anti-mutagenic properties

[52,53]. It is the most abundant from the HPLC analysis of the tomato powder mix for this study.

## CONCLUSION

The study revealed that the addition of ginger and *Hibiscus sabdariffa* to the tomato powder enhanced the nutritive components of the resultant powder mix. The phytochemical compounds contained in it also showed its inherent health-promoting effects and hence the possible use as a functional food that is cheap and readily available. When properly consumed, it may serve as a source of biologically active nutraceuticals for optimal health maintenance and preservation.

**Abbreviations:** AAS: Atomic-absorption spectrophotometer, GC-MS: Gas chromatography mass-

spectrometry HPLC: High-performance liquid chromatography, RT: retention time LOQ: Limit of quantification, LOD: Limit of detection.

**Authors' contributions:** The corresponding author substantially contributed to the conception and draft of the article. All others participated in data collection and revised the article critically for important intellectual content.

**Conflict of interest:** The authors declare there are no conflicts of interest.

**Acknowledgment/funding:** The authors declare no acknowledgments or funding.

## REFERENCES

- Mondal S, Soumya NPP, Mini S, Sivan SK: Bioactive compounds in functional food and their role as therapeutics. *Bioactive Compounds in Health and Disease* 2021, 4(3):24-39.  
DOI: <https://www.doi.org/10.31989/bchd.v4i3.786>
- Fiedor J, Burda K: Potential role of carotenoids as antioxidants in human health and disease. *Nutrients* 2014, 6(2):466-488.  
DOI: <https://www.doi.org/10.3390/nu6020466>
- Chaudhary P, Sharma A, Singh B, Nagpal AK: Bioactivities of phytochemicals present in tomato. *Journal of Food Science and Technology* 2018, 55(8):2833-2849.  
DOI: <https://www.doi.org/10.1007/s13197-018-3221-z>
- Wang C, Li M, Duan X, Abu-Izneid T, Rauf A, Khan Z, Mitra S, Emran TB, Aljohani AS, Alhumaydhi FA, Thiruvengadam M: Phytochemical and nutritional profiling of tomatoes: impact of processing on bioavailability-a comprehensive review. *Food Reviews International* 2022, July:1-25.  
DOI: <https://www.doi.org/10.1080/87559129.2022.2097692>
- Heuvelink E, (Editor): Tomatoes. 2<sup>nd</sup> edition. Boston: CABI; 2018, 27, 1-337. (Crop production science in horticulture series; Vol. 27).  
DOI: <https://www.doi.org/10.1079/9781780641935.0000>
- Kiralan M, Ketenoglu O: Utilization of tomato (*Solanum lycopersicum*) by-products: an overview. In *Mediterranean Fruits Bio-wastes: Chemistry, Functionality and Technological Applications*. Edited by Ramadan MF, Farag MA. Springer, Cham 2022:799-818.  
DOI: [https://www.doi.org/10.1007/978-3-030-84436-3\\_34](https://www.doi.org/10.1007/978-3-030-84436-3_34)
- Abu-Jdayil B, Banat F, Jumah R, Al-Asheh S, Hammad S: A comparative study of rheological characteristics of tomato paste and tomato powder solutions. *International Journal of Food Properties* 2004, 7(3):483-497.  
DOI: <https://www.doi.org/10.1081/JFP-200032940>
- Davoodi MG, Vijayanand P, Kulkarni S G, Ramana KVR: (2007). Effect of different pre-treatments and dehydration methods on quality characteristics and storage stability of tomato powder. *LWT-Food Science and Technology* 2007, 40(10):1832-1840.  
DOI: <https://www.doi.org/10.1016/j.lwt.2006.12.004>
- Bode A M, Dong Z: The amazing and mighty ginger. In *Herbal Medicine: Biomolecular and Clinical Aspects*. 2nd edition Edited by Benzie Wachtel-Galor S. Boca Raton (FL): CRC/Taylor and Francis; 2011: 131-156.
- Pattanittum P, Kunyanone N, Brown J, Sangkomkham US, Barnes J, Seyfoddin V, Marjoribanks J: Dietary supplements for dysmenorrhoea. *Cochrane Database of Systematic Reviews* 2016, 3:68-86.  
DOI: <https://www.doi.org/10.1002/14651858.CD002124.pub2>

11. Dev H, Sharma V: Genetic Variability in Ginger (*Zingiber officinale* Rosc.). *International Journal of Bio-resource and Stress Management* 2022, 13(7):709-717.  
DOI: <https://www.doi.org/10.23910/1.2022.2941a>
12. Hu W, Yu A, Wang S, Bai Q, Tang H, Yang B, Wang M, Kuang H: Extraction, Purification, Structural Characteristics, Biological Activities, and Applications of the Polysaccharides from *Zingiber officinale* Roscoe.(Ginger): A Review. *Molecules* 2023, 28(9):3855-3877.  
DOI: <https://www.doi.org/10.3390/molecules28093855>
13. Abidoye AO, Ojedokun FO, Fasogbon BM, Bamidele OP: Effects of sweet basil leaves (*Ocimum basilicum* L) addition on the chemical, antioxidant, and storage stability of roselle calyces (*Hibiscus sabdariffa*) drink. *Food Chemistry* 2022, 371:131170.  
DOI: <https://www.doi.org/10.1016/j.foodchem.2021.131170>
14. Da-Costa-Rocha I, Bonnlaender B, Sievers H, Pischel I, Heinrich M: *Hibiscus sabdariffa* L.—A phytochemical and pharmacological review. *Food Chemistry* 2014, 165:424-443.  
DOI: <https://www.doi.org/10.1016/j.foodchem.2014.05.002>
15. Riaz G, Chopra R: A review on phytochemistry and therapeutic uses of *Hibiscus sabdariffa* L. *Biomedicine and Pharmacotherapy* 2018, 102:575-586.  
DOI: <https://www.doi.org/10.1016/j.biopha.2018.03.023>
16. Kayode OT, Kayode AAA, Oyeboode TP. Chemometric analysis of ketogenic diet formulated from low-cost dietary fibers. *Functional Foods in Health and Disease* 2022; 12(1): 46-55.  
DOI: <https://www.doi.org/10.31989/ffhd.v12i1.857>
17. Sanusi J, Habsatu S, Abubakar N, Suleiman I, Musa DD: Comparative study of proximate and minerals composition of tomato cultivars in Sokoto, Sokoto State, Nigeria. *Fundamental Journal of Sciences* 2020, 4(4):409-414.  
DOI: <https://www.doi.org/10.33003/fjs-2020-0404-497>
18. Karpjuk UV, Al Azzam KM, Abudayeh ZHM, Kislichenko V, Naddaf, A., Cholak I, Yemelianova O: Qualitative and quantitative content determination of macro-minor elements in *Bryonia alba* L. roots using flame atomic absorption spectroscopy technique. *Advanced Pharmaceutical Bulletin* 2016, 6(2):285-291.  
DOI: <https://www.doi.org/10.15171/apb.2016.040>
19. Banu KS, Cathrine L: General techniques involved in phytochemical analysis. *International Journal of Advanced Research in Chemical Science* 2015, 2(4):25-32.
20. Harborne AJ: *Phytochemical methods: A guide to modern techniques of plant analysis*. 3<sup>rd</sup> edition. London: Chapman and Hall; 1998:33-80.  
DOI: <https://www.doi.org/10.1007/978-94-009-5921-7>
21. Adesanwo JK, Makinde O O, Obafemi CA: Phytochemical analysis and antioxidant activity of methanol extract and betulinic acid isolated from the roots of *Tetracera potatoria*. *Journal of Pharmacy Research* 2013,6(9):903-907.  
DOI: <https://www.doi.org/10.1016/j.jopr.2013.09.003>.
22. Evans, W. C., and Evans, D. (2009). Introduction. *Trease and Evans' Pharmacognosy*, 539. DOI: <https://doi.org/10.1016/b978-0-7020-2933-2.00055-1>
23. Anguelova T, Warthesen J: Lycopene stability in tomato powders. *Journal of Food Science* 2000, 65(1):67-70. DOI: <https://www.doi.org/10.1111/j.1365-2621.2000.tb15957.x>
24. Barba AO, Hurtado MC, Mata M S, Ruiz V F, De Tejada M L S: Application of a UV-vis detection-HPLC method for a rapid determination of lycopene and  $\beta$ -carotene in vegetables. *Food chemistry* 2006, 95(2):328-336.  
DOI: <https://www.doi.org/10.1016/j.foodchem.2005.02.028>
25. Tang N, An J, Deng W, Gao Y, Chen Z, Li Z: The metabolic and transcriptional regulatory mechanism associated with postharvest fruit ripening and senescence in cherry tomatoes. *Postharvest Biology and Technology* 2020, 168(11):1274:1-11.  
DOI: <https://www.doi.org/10.1016/j.postharvbio.2020.111274>
26. Aboagye-Nuamah F, Hussein YA, Ackun A: Biochemical properties of six varieties of tomato from Brong Ahafo region of Ghana as influenced by the ripening condition and drying. *African Journal of Food, Agriculture, Nutrition and Development* 2018, 18(1):13095-13109.  
DOI: <https://www.doi.org/10.18697/ajfand.81.16975>
27. Obadina A, Ibrahim J, Adekoya I: Influence of drying temperature and storage period on the quality of cherry and plum tomato powder. *Food Science & Nutrition* 2018, 6(4):1146-1153. DOI: <https://www.doi.org/10.1002/fsn3.658>.
28. Engin AS, Kuşçu A: The Effect of Different Drying Processes on Some Nutritional and Quality Criteria in Tomato Powder. *Gaziosmanpaşa Bilimsel Araştırma Dergisi* 2023, 12(1):194-204.
29. Karakaya S, Yılmaz N: Lycopene content and antioxidant activity of fresh and processed tomatoes and *in vitro* bioavailability of lycopene. *Journal of the Science of Food and Agriculture* 2007, 87(12):2342-2347. DOI: <https://www.doi.org/10.1002/jsfa.2998>
30. Degwale A, Asrat F, Eniyew K, Asres D, Tesfa T, Ayalew A: Influence of Dehydration Temperature and Time on Physicochemical Properties of Tomato (*Solanum lycopersicum* L.) Powder. *Frontiers in Sustainable Food Systems* 2022,

- 6:839385.  
DOI: <https://www.doi.org/10.3389/fsufs.2022.839385>
31. Nielsen SS: Introduction to food analysis. In Food analysis. 5<sup>th</sup> edition. Edited by Nielsen SS. Gaithersburg: Aspen Publishers; 1998:3-75.  
DOI: <https://www.doi.org/10.1007/978-3-319-45776-5>
  32. Mohammed SM, Abdurrahman AA, Attahiru M: Proximate analysis and total lycopene content of some tomato cultivars obtained from Kano State, Nigeria. *ChemSearch Journal* 2017, 8(1):64-69. DOI: <http://dx.doi.org/10.4314/cs.v8i1.9>
  33. Islary A, Sarmah J, Basumatary S: Proximate composition, mineral content, phytochemical analysis and *in vitro* antioxidant activities of a wild edible fruit *Grewia sapida* Roxb. ex DC. found in Assam of North-East India. *American Journal of Physiology, Biochemistry and Pharmacology* 2016, 5(1):1-11. DOI: <https://www.doi.org/10.5455/ajpb.20160422015354>
  34. Bello MI, Amubieya FO: Effects of Different Drying Methods on the Nutritional Status of *Lycopersicon Esculentum* (Tomato) Mill. *IOSR Journals of Biotechnology and Biochemistry* 2019, 5(6):69-71.  
DOI: <https://www.doi.org/10.9790/264X-0506016971>.
  35. Zoroddu MA, Aaseth J, Crisponi G, Medici S, Peana M, Nurchi VM: The essential metals for humans: a brief overview. *Journal of Inorganic Biochemistry* 2019, 195:120-129.  
DOI: <https://www.doi.org/10.1016/j.jinorgbio.2019.03.013>
  36. Aberoumand A, Deokule SS: Determination of elements profile of some wild edible plants. *Food Analytical Methods* 2009, 2(2):116-119.  
DOI: <https://www.doi.org/10.1007/s12161-008-9038-z>.
  37. Esmeeta A, Adhikary S, Dharshnaa V, Swarnamughi P, Maqsummiya ZU, Banerjee A, Pathak S, Duttaroy AK: Plant-derived bioactive compounds in colon cancer treatment: An updated review. *Biomedicine & Pharmacotherapy* 2022, 153(113384):1-12.  
DOI: <https://www.doi.org/10.1016/j.biopha.2022.113384>
  38. Fenni S, Hammou H, Astier J, Bonnet L, Karkeni E, Couturier C, Tourniaire F, Landrier JF: Lycopene and tomato powder supplementation similarly inhibit high-fat diet-induced obesity, inflammatory response, and associated metabolic disorders. *Molecular Nutrition and Food Research*, 2017, 61(9):1601083.  
DOI: <https://www.doi.org/10.1002/mnfr.201601083>
  39. Breinholt V, Lauridsen ST, Daneshvar B, Jakobsen J: Dose-response effects of lycopene on selected drug-metabolizing and antioxidant enzymes in the rat. *Cancer Letters* 2000, 154(2):201-210.  
DOI: [https://www.doi.org/10.1016/S0304-3835\(00\)00401-8](https://www.doi.org/10.1016/S0304-3835(00)00401-8)
  40. Hsu YM, Lai CH, Chang CY, Fan CT, Chen CT, Wu CH: Characterizing the lipid-lowering effects and antioxidant mechanisms of tomato paste. *Bioscience, Biotechnology and Biochemistry* 2008, 72(3):677-685. DOI: <https://www.doi.org/10.1271/bbb.70402>
  41. Böhm V: Lycopene and heart health. *Molecular Nutrition and Food Research* 2012, 56(2):296-303.  
DOI: <https://www.doi.org/10.1002/mnfr.769>
  42. Ayan HY, Artik NTD: *Production of sun-dried tomatoes (Lycopersitcum esculentum) and determination of changes during the process*. Ph.D. thesis. Ankara University Institute of Science, Food Engineering Department; 2010.
  43. Hussein JB, Sanusi MS, Filli KB: Evaluation of drying methods on the content of some bio-actives (lycopene, carotene and ascorbic acid) of tomato slices. *African Journal of Food Science* 2016, 10(12):359367.  
DOI: <https://www.doi.org/10.5897/AJFS2016.1470>
  44. Wong PK, Yusof S, Ghazali HM, Che Man YB: Physico-chemical characteristics of roselle (*Hibiscus sabdariffa* L.). *Nutrition & Food Science* 2002, 32(2):68-73.  
DOI: <https://www.doi.org/10.1108/00346650210416994>
  45. Yusop NM, Hishammudin MZ, Yahaya N, Ishak N, Ramlee NF, Zaini ZH: GC-MS screening of metabolites and antioxidant properties in four variants of *solanum lycopersicum* (tomato). *Journal of Engineering and Health Science* 2020, 4(1):61-82.
  46. Ramya S, Chandran M, King IJ, Jayakumararaj R, Loganathan T, Pandiarajan G, Kaliraj P, Pushpalatha GGL, Abraham GC, Vijaya V, Aruna D: Phytochemical Screening, GCMS and FTIR Profile of Bioactive Compounds in *Solanum lycopersicum* Wild Fruits collected from Palani Hill Ranges of the Western Ghats. *Journal of Drug Delivery and Therapeutic* 2022, 12(6):56-64.  
DOI: <https://www.doi.org/10.22270/jddt.v12i6.5665>
  47. Narain N, Galvao MDS, de Santana KL, da Silveira Moreira JDJ: Volatile compounds in tomato-based dried products. *Drying Technology* 2010, 28(2):232-239.  
DOI: <https://www.doi.org/10.1080/07373930903526905>
  48. Win DT: Oleic acid - The anti-breast cancer component in olive oil. Thailand. Assumption University Bangkok, faculty of Science and Technology: *AU J.T* 2005, 9(2):75-78. DOI: <https://www.thaiscience.info/journals/Article/AUJT/10290580.pdf>
  49. Teres S, Barceló-Coblijn G, Benet M, Alvarez R, Bressani R,

Halver JE, Escriba PV: Oleic acid content is responsible for the reduction in blood pressure induced by olive oil. *Proceedings of the National Academy of Sciences* 2008, 105(37):13811-13816.

DOI: <https://www.doi.org/10.1073/pnas.0807500105>

50. Sales-Campos H, Reis de Souza P, Crema Peghini B, Santana da Silva J, Ribeiro Cardoso C: An overview of the modulatory effects of oleic acid in health and disease. Mini-reviews in *Medicinal Chemistry* 2013, 13(2):201-210.

DOI: <https://doi.org/10.2174/138955713804805193>.

51. Mohamed R, Fernandez J, Pineda M, Aguilar M: Roselle (*Hibiscus sabdariffa*) seed oil is a rich source of  $\gamma$ -tocopherol. *Journal of Food Science* 2007, 72(3):S207-S211. DOI:

<https://www.doi.org/10.1111/j.1750-3841.2007.00285.x>

52. Ferrer A, Altabella T, Arró M, Boronat A: Emerging roles for conjugated sterols in plants. *Progress in Lipid Research* 2017, 67:27-37.

DOI: <https://www.doi.org/10.1016/j.plipres.2017.06.002>.

53. Kaur N, Chaudhary J, Jain A, Kishore L: Stigmasterol: a comprehensive review. *International Journal of Pharmaceutical Sciences and Research* 2011, 2(9):2259-2265.

ISSN: 0975-8232