



Identification of bioactive compounds in hydro-alcoholic extract of cinnamon by GC-MS analysis and evaluation of its in-vitro anti-gastric cancer potential

Mina Owrang¹, Alam Ara Gholami¹, Alireza Rafiei², Maryam Talei^{1*}

¹Faculty of Medicine, Sari branch, Islamic Azad University, Sari, Iran; ² Department of Immunology, Molecular and Cell Biology Research Center, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran.

*Corresponding Author: Mina Owrang, Faculty of Medicine, Sari branch, Islamic Azad University, Sari, Iran.

Submission Date: February 27th, 2024; **Acceptance Date:** May 23rd, 2024; **Publication Date:** May 27th, 2024

Please cite this article as: Owrang M., Gholami A. A., Rafiei A., Talei M. Identification of bioactive compounds in hydro-alcoholic extract of cinnamon by GC-MS analysis and evaluation of its in-vitro anti-gastric cancer potential. *Bioactive Compounds in Health and Disease* 2024; 7(5): 265-273. DOI: <https://www.doi.org/10.31989/bchd.v7i5.1286>

ABSTRACT

Background: Despite the availability of various treatment methods for gastric cancer, herbal medicine remains a promising candidate. This study aimed to identify the bioactive compounds in the hydroalcoholic extract (CHE) of cinnamon (*Cinnamomum zeylanicum*) using gas chromatography-mass spectrometry (GC-MS) analysis.


Objectives: To study the phytochemical constituents from the hydroalcoholic extract (CHE) of cinnamon (*Cinnamomum zeylanicum*).

Methods: The CHE was tested for cytotoxicity against a normal human gastric epithelium cell line (GES-1) and a human gastric adenocarcinoma cell line (AGS) as a model of gastric cancer.

Results: The analysis revealed six main phytochemical compounds in the CHE, with cinnamaldehyde being the predominant compound. The CHE showed potent cytotoxicity against the AGS cell line at concentrations ranging from 100-600 µg/mL, significantly reducing the viability of these cancerous cells to 20.88% compared to the control group (p<0.05). In contrast, the CHE did not significantly affect the viability of GES-1 cells.

Conclusion: The findings obtained suggest a promising avenue for finding new anticancer compounds as potent chemotherapeutic agents for gastric cancer.

Keywords: anticancer; bioactive compounds; cinnamon; cytotoxicity



INTRODUCTION

Gastric cancer (GC), also known as stomach cancer, is a dangerous disease that leads to the death of numerous people worldwide. According to the Global Cancer Observatory (GCO) reports, GC is the fifth most common cancer and, following lung and colorectal cancer, it is considered the third deadliest cancer, with a 5-year survival rate of less than 25% and accounting for 1 in 12 cancer-related deaths globally [1]. Therefore, GC is a severe public health concern [2]. The main risk factors for GC worldwide include *Helicobacter pylori* infection, smoking, and high salt intake [3]. Currently, common treatments for GC include gastrectomy, chemotherapy, and radiation therapy [4]. However, these methods are limited by their side effects or toxicity in GC patients [5]. In addition, resistance to anticancer drugs following chemotherapy further restricts the success of treatment [6]. Consequently, finding more effective and less toxic therapies for managing GC has become a significant focus of recent research.

One of the main candidates for cancer therapy is the use of herbal medicine, which offers a safe, non-toxic,

and readily available source of treatment [7]. According to the Functional Food Center (FFC), functional foods (FFs) are foods obtained from nature or derived from food processing that contain a suitable amount of biologically active compounds; these compounds provide health benefits and reduce the risk or symptoms of diseases [8]. Functional foods and bioactive compounds can be helpful in treating various clinical problems, including obesity [9], oxidative stress [10], inflammation [11-12], blood cholesterol [13], and so on [14-15]. Many studies have demonstrated that natural compounds in medicinal herbs can reduce the risk of GC [16-17]. The therapeutic effects of these herbs are primarily attributed to their phytochemicals [18]. Previous research has identified several mechanisms by which natural compounds in medicinal herbs exert positive effects on GC, including inhibiting cell proliferation and *Helicobacter pylori* infection, inducing apoptosis and autophagy, suppressing cell metastasis, and exhibiting anti-angiogenic properties [19-21].

Thus, it is important to focus on the phytochemicals in the present study. Among various herbs, cinnamon

holds a special place in traditional medicine and has been used since ancient times for its health-promoting potential [22]. Ariaee-Nasab et al. (2014) evaluated the anticancer potential of a water extract of cinnamon on human cancer cell lines (AGS, HeLa, MCF-7 and MDA-MB-234), demonstrating its anti-neoplastic activity [23]. Despite its anticancer potential, limited studies focus on the interaction of cinnamon with the human gastric adenocarcinoma cell line (AGS). In the present study, we determined the phytochemical constituents from the hydroalcoholic extract (CHE) of cinnamon (*Cinnamomum zeylanicum*) using gas chromatography-mass spectrometry (GC-MS). The cytotoxicity of the cinnamon extraction was assayed in the normal human gastric epithelium cell line (GES-1) and human gastric adenocarcinoma cell line (AGS) as a model of gastric cancer.

METHODS

Cinnamon hydro-alcoholic extraction: The method described by Anderson et al. was used to prepare the cinnamon hydro-alcoholic extract (CHE) [24]. Cinnamon powder was obtained from cinnamon bark imported from China and sold in Iran. To prepare the CHE, 200 g of cinnamon powder was dissolved in 500 mL of 0.1 N acetic acid and boiled for 20 min. The mixture was then centrifuged, and the supernatant was mixed with absolute ethanol at a ratio of 1:4. The resulting mixture was kept overnight at 4 °C, filtered using Whatman No. 1 filter paper, and dried in an oven at 60°C.

Gas chromatography-mass spectrometry (GC-MS): A gas chromatography-mass spectrometry (GC-MS) analysis of the CHE was conducted using an Agilent 6890 N system (US), following established protocols [25]. The GC-MS system was equipped with an HP 5989 mass spectrometer detector. The Agilent 5MS-HP column (US)

used was 30 m in length, with an inner diameter of 0.025 mm and a particle size of 0.25 µm. Nitrogen was used as a carrier gas to transport the evaporated extract into the column. For data analysis, Agilent G1701DA MSD ChemStation software (US) was utilized.

Cell culture: The human gastric epithelial cell line (GES-1) and the human gastric adenocarcinoma cell line (AGS) were obtained from the National Cell Bank of Iran (Tehran, Iran). The GES-1 and AGS cells were cultured in H-DMEM (HyClone) and RPMI 1640 (Gibco) media in 96-well (TC) plates (NUNC). The media were supplemented with 10% fetal bovine serum (Invitrogen) and penicillin–streptomycin (Sigma). The cells were maintained at 37°C in a 95% humidified atmosphere enriched by 5% CO₂.

Cell exposure and cell cytotoxicity assay: The resulting powder, or the CHE, was dissolved in dimethyl sulfoxide (DMSO) to prepare a stock solution with a concentration of 100 mg/mL. An MTT assay was used to evaluate the cytotoxic effects of CHE on GES-1 and AGS cells. Cells were seeded in 96-well plates at a density of 1×10^4 cells per well and incubated for 24 h. The plates were then treated with various concentrations of CHE (0, 100, 200, 400, 600, 800, and 1000 µg/mL) and incubated for 24, 48, and 72 h. After incubation, the medium in each well was removed and replaced with 100 µL of MTT at 0.5 mg/mL concentration. The plates were incubated for an additional 4 h at 37°C in a humidified atmosphere with 5% CO₂. Following this, the MTT solution was removed, and 100 µL of DMSO solution was added to each well to dissolve the purple formazan crystals. Optical absorbance was measured using an Awareness Technologies Stat Fax 2100 ELISA reader (US) at a wavelength of 570 nm. The results were reported as the IC₅₀ values and cell viability percentages [26].

Statistical analysis: The statistical analysis was performed using OriginPro 2018 9.5 version Software. The results were analyzed using one-way ANOVA and Tukey's multiple-range test to compare one factor among multiple groups. Data are presented as the mean \pm SD at a 95% confidence level. The probit analysis method was used to transform the growth inhibition data, and the IC₅₀ values were derived by linear regression from plots of probit units against the correspondent log concentrations.

RESULTS:

Cell cytotoxicity: Treatment with different concen-

trations of CHE did not exhibit high cytotoxicity in GES-1 cells, except at concentrations of 800 and 1000 μ g/mL, which led to a significant decrease in the percentage of viable cells ($p < 0.05$) (Figure 2).

In contrast, the survival of AGS cells showed no significant change after treatment with 100 and 200 μ g/mL of CHE ($p > 0.05$) (Figure 2). However, treatment with CHE at concentrations ranging from 400-1000 μ g/mL resulted in a significant decrease in cell viability compared to the control group ($p < 0.05$) (Figure 2). IC₅₀ values for AGS cells at 24, 48, and 72 h are presented in Table 1, with the lowest IC₅₀ value observed after 72 h of CHE treatment.




Figure 1. GC/MS spectrogram for cinnamon hydro-alcoholic extraction.

Table 1. Phytocomponents identified in Cinnamon hydro-alcoholic extraction.

| Pike number | RT | Name of the compound | Peak Area % | RF | CAS |
|-------------|--------|---|-------------|--------|-------------|
| 1 | 21.053 | 2-Propenal, 3-phenyl, 3-phenylpropenal cinnamal | 21.47 | 39272 | 000104-55-2 |
| 1 | 21.053 | 2-Propenal, 3-phenyl-(CAS), cinnamaldehyde 3-phenylpropenal cinnamal | 21.7 | 39276 | 2-55-000104 |
| 1 | 21.053 | 2-Propenal, 3-phenyl-cinnamaldehyde acrolein, 3-phenyl-\$\$ cinnamal | 21.7 | 39274 | 2-55-000104 |
| 7 | 26.523 | 1S, cis-calamenene, 6-dimethyl-1,2,3,4-tetrahydronaph thalene | 8.64 | 86283 | 2-77-000483 |
| 17 | 48.169 | Megastigma-5(11),7,9-trien-1H-purin-2-amine, 6-methoxy-N-meth 1,2-dihydro-4-(2-propenylamino) | 7.53 | 159760 | 072523569 |
| 6 | 25.808 | alpha-Muurolene | 6.11 | 191751 | 9-22031983 |




Figure 2. Effect of cinnamon hydro-alcoholic extract with various concentration on cell viability of GES-1 (a) and AGS (b) cell line at 24, 48, and 72 h. Different letters in each group indicate significant differences, P < 0.05. (n = 3, Mean ± SD).

Table 2. The value of test solution IC₅₀ on gastric cancer cell AGS using Probit analysis. (IC₅₀ ± SD).

| IC ₅₀ value in AGS cells (µg/mL) | Time |
|---|----------|
| 940 ± 50 | 24 hours |
| 740 ± 35 | 48 hours |
| 490 ± 22 | 72 hours |

DISCUSSION

The use of natural products as potential anticancer agents holds great promise in current therapeutic strategies. Moreover, structural modifications of natural compounds can help generate novel clinical candidates targeting cancer cells [27]. Cinnamon, known for its antitumor properties demonstrated in previous studies [28], stands out among similar herbs. Hence, this study selected cinnamon as the focal plant for investigation.

The present study used the hydro-alcoholic method to extract cinnamon. This method offers several advantages, notably its environmental friendliness, safety, and efficiency. Moreover, this method can be used to extract polar compounds, such as cinnamaldehyde [29]. A previous study showed the effectiveness of the hydro-alcoholic method in reducing the proliferation and growth of cancer cells, as observed

with the plant *Capparis spinosa*, attributed to the presence of flavonoids, including cinnamaldehyde [30].

In this study, results showed that the main compound in CHE was cinnamaldehyde, a finding consistent with previous research. Uma et al. similarly reported cinnamaldehyde as the dominant compound in the alcoholic extract of *Cinnamon zeylanicum* [31]. Previous studies have revealed that aromatic aldehydes, like cinnamaldehyde, have unique biological and pharmacological properties, including antioxidant, antimutagenic, and anti-inflammatory activities [32-33]. Recently, cinnamaldehyde has gained considerable attention as a powerful anticancer agent [33]. It shows efficacy in treating various conditions, such as diabetes, atherosclerosis, cancer, inflammation, and cardiovascular disease. Its anticancer potential originates from its ability to induce apoptosis and cell cycle arrest. Moreover, it also has an inhibitory role in angiogenesis, oxidative stress, inflammation, cellular invasion, and metastasis [34].

In the present study, cytotoxicity investigations revealed that CHE treatment did not cause significant changes in viability and proliferation of GES-1 cells, while it exerted a strong inhibitory effect on the viability and proliferation of gastric cancer cells (AGS cell line). In one study, cinnamaldehyde extracted from cinnamon in hepatoma cells (HepG2 and Hep3B) inhibited cell proliferation [35]. In another study, polyphenols containing cinnamaldehyde inhibited cell proliferation and induced cell death in human ovarian cancer cells via cell cycle arrest [36]. Furthermore, low molar concentrations of cinnamon extract have been found to inhibit cell proliferation in the melanoma A375 cell line by increasing intracellular reactive oxygen species (ROS) [27]. Consistent with the observations of the present study, previous research has reported that cinnamaldehyde derived from cinnamon extract exhibits a potent cytotoxic effect on cancer cells, while

demonstrating minimal cytotoxicity to normal cells [32-37].

The present study also observed that treatment with CHE at concentrations of 800 and 1000 µg/mL did not induce significant changes in the viability and proliferation of GES-1 normal cells. However, CHE had a strong inhibitory effect on the proliferation and viability of gastric cancer cells. In a past study, it was demonstrated that trans-cinnamaldehyde, not CHE, at a concentration of 2000 µg/mL reduced the development of AGS cells [38]. Trans-cinnamaldehyde was found to induce apoptosis in the AGS cell line by increasing the levels of cleaved caspase-9, cleaved PARP, p53, and Bax proteins [34].

Although studies on the effect of cinnamon extract on gastric cancer and its underlying molecular mechanisms are still in their infancy, present evidence suggests that cinnamaldehyde, a prominent component of cinnamon, may suppress gastric cancer cell growth via apoptotic agents, such as caspase-3 cleavage and Bcl-2 reduction, along with autophagic markers.

CONCLUSION

The results of the present study supported the idea that the hydroalcoholic extract of cinnamon, which contains cinnamaldehyde, showed high anticancer properties in gastric cancer cells, while demonstrating no toxic effect on healthy gastric cells. These findings warrant further investigation to explore the anti-gastric cancer effects of HCE in vivo and to evaluate its therapeutic potential against gastric cancer in clinical trials.

List of Abbreviations: CHE, cinnamon hydroalcoholic extract; GES-1, human gastric epithelium cell line; AGS, human gastric adenocarcinoma cell line; GC, gastric cancer; GCO, Global Cancer Observatory; DMSO, dissolved in dimethyl sulfoxide.

Authors' Contributions: M. O. designed the study and wrote the manuscript, A. A. G. edited the manuscript, A. R. performed the experiments, and M. T. performed the statistical analysis.

Competing Interests: The authors have no conflict of interest.

Acknowledgement and Funding: The authors would like to thank Islamic Azad University for providing instruments.

REFERENCES:

1. Yamaoka Y: How to eliminate gastric cancer-related death worldwide? *Nat Rev Clin Oncol*, 2018; 15(7): 407-408. DOI: <https://doi.org/10.1038/s41571-018-0029-8>
2. Yan S, Li B, Bai Z, Wu JQ, Xie DW, Ma YC, Ma XX, Z JH, Guo XJ: Clinical epidemiology of gastric cancer in Hehuang valley of China: A 10-year epidemiological study of gastric cancer. *World J Gastroenterol*, 2014; 20(30): 10486-10494. DOI: <https://doi.org/10.3748/wjg.v20.i30.10486>
3. Farmanfarma KK, Mahdavifar N, Hassanipour S, Salehiniya H: Epidemiologic study of gastric cancer in Iran: A systematic review. *Clin Exp Gastroenterol*, 2020; 13: 511-542. DOI: <https://doi.org/10.2147/CEG.S256627>
4. Izuishi K, H Mori: Recent Strategies for Treating Stage IV Gastric Cancer: Roles of Palliative Gastrectomy, Chemotherapy, and Radiotherapy. *J Gastrointest Liver Dis*, 2016; 25(1): 87-94. DOI: <https://doi.org/10.15403/jgld.2014.1121.251.rv2>
5. Ren F, Li S, Zhang Y, Zhao Z, Wang H, Cui Y: Efficacy and safety of intensity-modulated radiation therapy versus three-dimensional conformal radiation treatment for patients with gastric cancer: A systematic review and meta-analysis. *Radiat Oncol*, 2019; 14(1): 1-9. DOI: <https://doi.org/10.1186/s13014-019-1294-0>
6. Yu LL, Wu JG, Dai N, Yu HG, Si JM: Curcumin reverses chemoresistance of human gastric cancer cells by downregulating the NF- κ B transcription factor. *Oncol Rep*, 2011; 26(5): 1197-1203. DOI: <https://doi.org/10.3892/or.2011.1410>
7. Ahmad S, Munir S, Zeb N, Ullah A, Khan B, Ali J, Bilal M, Omer M, Alamzeb M, Salman SM, Ali S. Green nanotechnology: A review on green synthesis of silver nanoparticles—an ecofriendly approach. *Int J Nanomedicine*, 2019; 14: 5087-5107. DOI: <https://doi.org/10.2147/IJN.S200254>
8. Martirosyan D, Kanya H, and Nadalet C: Can functional foods reduce the risk of disease? Advancement of functional food definition and steps to create functional food products: *Functional Foods in Health and Disease*, 2021; 11(5): 213-221. DOI: <https://doi.org/10.31989/ffhd.v11i5.788>
9. Papagianni O, Bountziouka V, Maniati E, Pegkos A, Dimou C, Karantonis HC, Koutelidakis AE: The frequency of selective functional foods consumption is related to improved anthropometric obesity indices: A cross-sectional study in Greek and Cyprian adults. *Functional Food Science*, 2024; 4(4): 134-152. DOI: <https://doi.org/10.31989/ffs.v4i4.1274>
10. Kim YJ, Jung KY, Lee JW, Om AS: Protective effects of Quercus salicina blume leaves aqueous extracts against cadmium-induced oxidative stress and cell cytotoxicity in HK-2 cells. *Functional Foods in Health and Disease*, 2023; 13(11): 648-663. DOI: <https://doi.org/10.31989/ffhd.v13i11.1057>
11. Moghrovyan A, Ginovyan M, Avtandilyan N, Parseghyan L, Voskanyan A, Sahakyan N, Darbinyan A: The possible anti-inflammatory properties of hydro-ethanolic extract of Oregano. *Functional Foods in Health and Disease*, 2023; 13(10): 476-486. DOI: <https://doi.org/10.31989/ffhd.v13i10.1211>
12. Sholmire A, Ishii T, Leischner L, Garland B, Yamauchi Y, Shirako S, Nishidono Y, Ikeya Y, Corey L, Nishizawa M. Fatty acids in Plantago asiatica seeds are responsible for the production of the pro-inflammatory mediator nitric oxide. *Bioactive Compounds in Health and Disease*, 2024; 7(5): 233-244. DOI: <https://doi.org/10.31989/bchd.v7i5.1337>
13. Dehghani P, Masjedi M, Shariati L, Vaseghi G, Dana N, Zeinalian M, Asgary S. Exploring the potential of pomegranate (Punica granatum) peel extract and punicalagin as novel anti-hypercholesterolemic agents. *Functional Foods in Health and Disease*, 2024; 14(4): 270-281. DOI: <https://doi.org/10.31989/ffhd.v14i4.1320>
14. Salman I, Martirosyan D: Migraine management: A review of healthy diets and bioactive compounds. *Functional Food Science*, 2023; 3(8): 129-153. DOI: <https://doi.org/10.31989/ffs.v3i8.1149>
15. Grigoryan B, Mikayelyan M: The investigation of bioactive compounds in the Charentsi grape variety and its derived wines. *Bioactive Compounds in Health and Disease*, 2023; 6(11): 303-314. DOI: <https://doi.org/10.31989/bchd.v6i11.1170>

16. Zaidi SF, Muhammad JS, Usmanghani K, Sugiyama T: Pharmacological ins and outs of medicinal plants against *Helicobacter pylori*: A review. *Pak J Pharm Sci*, 2015; 28(3): 1171-1176.
17. Huang WJ, Ruan S, Wen F, Lu Xn, Gu Sp, Chen Xx, Liu M, Shu P: Multidrug resistance of gastric cancer: The mechanisms and Chinese medicine reversal agents. *Cancer Manag Res*, 2020; 12: 12385-12394.
DOI: <https://doi.org/10.2147/CMAR.S274599>
18. Wada K, Tsuji M, Tamura T, Konishi K, Kawachi T, Hori A, Tanabashi S, Matsushita S, Tokimitsu N, Nagata C: Soy isoflavone intake and stomach cancer risk in Japan: From the Takayama study. *Int J Cancer*, 2015; 137(4): 885-892.
DOI: <https://doi.org/10.1002/ijc.29437>
19. Chen W, Zhao Z, Li Y: Simultaneous increase of mycelial biomass and intracellular polysaccharide from *Fomes fomentarius* and its biological function of gastric cancer intervention. *Carbohydr Polym*, 2011; 85(2): 369-375.
DOI: <https://doi.org/10.1016/j.carbpol.2011.02.035>
20. Kim TW, Lee SY, Kim M, Cheon C, Ko SG: Kaempferol induces autophagic cell death via IRE1-JNK-CHOP pathway and inhibition of G9a in gastric cancer cells. *Cell Death Dis*, 2018; 9(9): 875.
DOI: <https://doi.org/10.1038/s41419-018-0930-1>
21. Mahady GB, Pendland SL, Yun G, Lu ZZ: Turmeric (*Curcuma longa*) and curcumin inhibit the growth of *Helicobacter pylori*, a group 1 carcinogen. *Anticancer Res*, 2002; 22(6C): 4179-4181. DOI: <https://doi.org/10.1021/jf034916b>
22. Dutta A, Chakraborty A: Cinnamon in anticancer armamentarium: A molecular approach. *J Toxicology*, 2018; 8978731. DOI: <https://doi.org/10.1155/2018/8978731>
23. Ariaee-Nasab N, Vahedi Z, Vahedi F: Inhibitory effects of cinnamon-water extract on human tumor cell lines. *Asian Pac J Trop Dis*, 2014; 4(2), S975-S978.
DOI: [https://doi.org/10.1016/S2222-1808\(14\)60768-4](https://doi.org/10.1016/S2222-1808(14)60768-4)
24. Anderson RA, Broadhurst CL, Polansky MM, Schmidt WF, Khan A, Flanagan VP, Schoene NW, Graves DJ: Isolation and characterization of polyphenol type-A polymers from cinnamon with insulin-like biological activity. *J Agric Food Chem*, 2004; 52(1): 65-70.
DOI: <https://doi.org/10.1021/jf034916b>
25. Emami S, Ahmadi M, Nasiraie LR, Shahidi SA, Jafarizadeh-Malmiri H: Cinnamon extract and its essential oil nanoliposomes—preparation, characterization and bactericidal activity assessment. *Biologia*, 2022; 77(10): 3015-3025.
DOI: <https://doi.org/10.1007/s11756-022-01164-x>
26. Plumb JA: Cell sensitivity assays: the MTT assay. *Methods Mol Med*, 1999; 28: 25-30.
DOI: <https://doi.org/10.1385/1-59259-687-8:25>
27. Chang S, Qin D, Wang L, Zhang M, Yan R, Zhao C: Preparation of novel cinnamaldehyde derivative–BSA nanoparticles with high stability, good cell penetrating ability, and promising anticancer activity. *Coll Surf A Colloid Surf A Physiochem Eng Asp*, 2021; 624: 126765.
DOI: <https://doi.org/10.1016/j.colsurfa.2021.126765>
28. Sadeghi S, Davoodvandi A, Pourhanifeh MH, Sharifi N, ArefNezhad R, Sahebnaasagh R, Moghadam SA, Sahebkar A, Mirzaei H: Anti-cancer effects of cinnamon: Insights into its apoptosis effects. *Eur J Med Chem*, 2019; 178: 131-140.
DOI: <https://doi.org/10.1016/j.ejmech.2019.05.067>
29. Ghazizadeh J, Hamedeyazdan S, Torbati M, Farajdokht F, Fakhari A, Mahmoudi J, Araj-Khodaie M, Sadigh-Eteghad S: *Melissa officinalis* L. hydro-alcoholic extract inhibits anxiety and depression through prevention of central oxidative stress and apoptosis. *Exp Physiol*, 2020; 105(4): 707-720.
DOI: <https://doi.org/10.1113/EP088254>
30. Moghadamnia Y, Kani SNM, Ghasemi-Kasman M, Kani MTK, Kazemi S: The anti-cancer effects of *Capparis spinosa* hydroalcoholic extract. *Avicenna J Med Biotechnol*, 2019; 11(1): 43-47.
31. Uma B, Prabhakar K, Rajendran S, Lakshmi SY: Studies on GC/MS spectroscopic analysis of some bioactive antimicrobial compounds from *Cinnamomum zeylanicum*. *J Med Plant Res*, 2009; 8(31): 125-131.
32. Ka H, Park HJ, Jung HJ, Choi JW, Cho KS, Ha J, Lee KT: Cinnamaldehyde induces apoptosis by ROS-mediated mitochondrial permeability transition in human promyelocytic leukemia HL-60 cells. *Cancer Lett*, 2003; 196(2): 143-152.
DOI: [https://doi.org/10.1016/S0304-3835\(03\)00238-6](https://doi.org/10.1016/S0304-3835(03)00238-6)
33. Kim TW: Cinnamaldehyde induces autophagy-mediated cell death through ER stress and epigenetic modification in gastric cancer cells. *Acta Pharmacol Sin*, 2022; 43(3): 712-723. DOI: <https://doi.org/10.1038/s41401-021-00672-x>
34. Banerjee S, Banerjee S: Anticancer potential and molecular mechanisms of cinnamaldehyde and its congeners present in the cinnamon plant. *Physiol*, 2023; 3(2): 173-207.
DOI: <https://doi.org/10.3390/physiologia3020013>
35. Chuang LY, Guh JY, Chao LK, Lu YC, Hwang JY, Yang YL, Cheng TH, Yang WY, Chien YJ, Huang JS: Anti-proliferative effects of cinnamaldehyde on human hepatoma cell lines. *Food Chem*, 2012; 133(4): 1603-1610.
DOI: <https://doi.org/10.1016/j.foodchem.2012.02.059>

36. Shin SY, Jung H, Ahn S, Hwang D, Yoon H, Hyun J, Yong Y, Cho HJ, Koh D, Lee, YH, Lim Y: Polyphenols bearing cinnamaldehyde scaffold showing cell growth inhibitory effects on the cisplatin-resistant A2780/Cis ovarian cancer cells. *Bioorg Med Chem*, 2014; 22(6): 1809-1820. DOI: <https://doi.org/10.1016/j.bmc.2014.01.058>
37. Han DC, Lee MY, Shin KD, Jeon SB, Kim JM, Son KH Kim HC, Kim HM, Kwon BM: 2'-Benzoyloxycinnamaldehyde induces apoptosis in human carcinoma via reactive oxygen species. *J Biol Chem*, 2004; 279(8): 6911-6920. DOI: <https://doi.org/10.1074/jbc.M309708200>
38. Milani AT, Rashidi S, Mahmoudi R, Douna BK: Cytotoxic activity of epigallocatechin and trans-cinnamaldehyde in gastric cancer cell line. *Asian Pac J Cancer Biol*, 2019; 4(4): 71-74. DOI: <https://doi.org/10.31557/apicb.2019.4.4.71-74>